



**Dr. Sarah Miller**  
MD, FRCS

1636 College Avenue | Regina, SK. | S4P 1B6  
T. 306-546-5504 | F. 306-205-8939 | drsarahmiller.ca

## Gastroscopy Instructions

You have been booked for a gastroscopy with Dr. Miller on \_\_\_\_\_ at the \_\_\_\_\_ . Please present to the admitting department at \_\_\_\_\_ and then proceed to the endoscopy unit, which is a scent-free environment.

### Gastroscopy Checklist

- Upon receipt of this notice, please call (306) 546-5504 to confirm your appointment. Any **rebooking or cancellations must be made 72 hours prior** given lengthy waitlists. If not, there may be a charge associated with not cancelling in this timeframe (exceptions made for extenuating circumstances).
- Please notify Dr. Miller's office if you are taking any **blood thinners** - eg) Warfarin, Clopidogrel/Plavix, Dabigatran/Pradaxa, Rivaroxaban/Xarelto, Apixaban/Eliquis. Depending on your medical history, these may need to be stopped prior to your procedure. Please also notify the office if you have a pacemaker or ICD. If you are taking 81 mg of aspirin, this may be continued.
- Stop eating any solid foods 8 hours prior to your procedure.**
  - During this time, follow a **clear fluid diet**. A **clear fluid diet** includes water, clear broth, clear fruit juices, Jell-o, popsicles without pulp, carbonated beverages, sports drinks, and tea and coffee (black or with sugar). Do not eat or drink any dairy products (e.g. milk, cream, yogurt, cheese, ice cream). Avoid any red-coloured beverages.
  - Stop drinking any clear liquids at least **2 hours before** your scheduled admitting time.
  - Do not consume any alcohol **24 hours** prior to your procedure.

### The Day of your Gastroscopy

- Medical considerations:
  - If you have diabetes, **do not take your short acting insulin or oral diabetic medications the morning** of your procedure. Remember to use glucose-containing clear liquids to help you maintain your blood sugar. If you need help managing this prior to your procedure, please contact your family physician.
  - All other medications may be taken on the morning of your procedure as usual (unless instructed otherwise). Please bring your list to the hospital.
  - If you have sleep apnea and use a CPAP machine, please bring this to your appointment.
- Plan to be at the hospital for approximately 3-4 hours. Leave any jewelry or valuables at home. **You must have someone to pick you up following the procedure if you receive sedation.** You cannot drive for 24 hours following administration of sedative medications. If you choose to complete the gastroscopy without sedation, you may drive yourself home afterwards.